



Anterior Labrum Repair Protocol

Stage I (0-4 weeks):

Key Goals:

- Protect the newly repaired shoulder.
- Allow for decreased inflammation and healing.
- Maintain elbow, wrist and hand function.
- Maintain scapular control.
- Begin passive abduction and forward flexion

1. Immobilizer use:

- a. The immobilizer will be placed on patient's shoulder in surgery.
- b. The patient may remove the immobilizer for dressing and hygiene.
- c. The patient should wear the immobilizer full-time for four weeks.

2. Restrictions:

- a. No shoulder external rotation.
 - i. The capsular repair is stressed with movement into external rotation. Since the repair is performed with the shoulder in a neutral position external rotation must be limited for six weeks following the repair.
- b. When arm is out of the immobilizer, forearm must be touching abdomen.
- c. Acceleration of rehabilitation for "fast healers" may reduce results and lead to long-term problems.

3. Exercises:

- a. Pendulum exercises.
- b. May begin passive shoulder range of motion beginning the week after surgery
 - i. Passive external rotation to neutral position only for weeks 1 - 4
- c. Active assistive range of motion of the involved elbow, wrist and hand in the plane of the body. The patient may progress to active range of motion as comfort improves.
- d. Scapular control exercises (Immobilizer on)
- e. Core training (Immobilizer on)

Stage II (5-15 weeks):

Key Goals:

- Full active elevation at 12 weeks from surgery.
- Surgical shoulder external rotation of 80% of uninvolved shoulder.
- Normal scapular mechanics 12 weeks from surgery.
 - Scapular mechanics should be evaluated on a regular basis.
- Normal scapular stabilizer, rotator cuff and core strength at 16 weeks from surgery.

1. Weeks 5-6:

- a. Brace use:
 - i. Immobilizer will be used at this time while sleeping until six weeks post-op.
 - ii. Sling is worn during the day for comfort. Wean as comfort improves.
- b. Range of motion:
 - i. External rotation:
 1. Passive to active assistive to active range of motion as able.
 - 2. Limited to 20 degrees maximum until 6 weeks from surgery.**
 - 3. No subscapularis or anterior shoulder stretching until 6 weeks from surgery.**
 - ii. Internal rotation:
 1. Passive to active assistive to active range of motion as able.
 - a. Begin in supine with scapula stabilized, and progress to other postures as tolerated.
 - iii. Flexion/Scaption/Abduction:
 1. Passive to active assistive to active range of motion as able.
 - a. Supine with scapula stabilized.
 - iv. Gleno-humeral mobilizations:
 1. No anterior glides until 10 weeks from surgical date.
- c. Balance training:
- d. Strengthening (4 weeks):
 - i. Isometric strengthening:
 - a. Internal/external rotation:

1. **If open surgical procedure, NO internal rotation strengthening until six weeks post-op.**

ii. Core training:

2. Week 7:

- a. Immobilizer use at night can be discontinued.
- b. Range of motion:
 1. As tolerated no limits.
- c. Strengthening:
 - i. Scapular stabilizer strengthening:
 - ii. Core training:

3. Week 8:

- **Warning: No soreness with the above rotator cuff strengthening.**
 - **The program must be modified to avoid cuff aggravation.**
- a. Balance training:
 - b. Range of motion:
 - i. No anterior apprehension or impingement.
 - ii. **Scapular mechanics need to be functioning properly and if not need to be addressed.**
 - iii. Hip mobility:
 - c. Strengthening:
 - i. Scapular mechanics:
 - ii. Forearm strengthening:
 - iii. Rotator cuff strengthening:
 - iv. Core training

4. Week 12:

- a. Testing:
 - i. Full pain free active range of motion for elevation and internal rotation.

- ii. A 20 degree difference in shoulder external rotation is acceptable.
- iii. Normal scapular mechanics.
- iv. ROM is within 10 degrees of other side.
 - 1. ROM should be within 5 degrees or less by 16 weeks.
- v. Int Rotation difference is less than 20 degrees or 2 spinal levels.
- vi. Squat screen (FMS):
- vii. Hurdle step screen (FMS):
- viii. Shoulder mobility screen (FMS):
- ix. Hand held dynamometer:
 - 1. 0 degrees with arm at side IR and ER.
 - 2. Seated IR and ER at 90 degrees of abduction and 45 degrees of external rotation.
 - 3. ER/IR=65%

Warning:

- **Any deficits in mobility, stability, or scapular mechanics need to be addressed now prior to beginning return to throw program at 20 weeks.**

- b. Range of motion:
 - i. Any flexibility deficits need to be addressed before return to program begins at 16 weeks.
 - 1. See above testing.
 - 2. Begin external rotation/pectoral stretching.**
- c. Strengthening:
 - i. Scapular stabilizer:
 - ii. Rotator cuff:
 - iii. Plyometric training
 - 1. Upper extremity.
 - 2. Lower extremity.
 - iv. Core training:
 - v. Endurance training:

Stage III (Weeks 20-26)

Initiation of Interval Sport Program for Baseball, Tennis, and Golf:

- Return-to-sport activities after injury that include attention to the entire body.
 - A gradual progression of applied forces to lessen the chance of re-injury.
 - Proper warm-up and maintenance exercises.
 - Proper biomechanics to minimize the chance of re-injury.
 - Variability is based on each athlete's skill, level, goals and injury.
 - Program needs to be followed rigidly. Some athletes will try and rush through the plan.
 - No skipping of steps is allowed.
 - Patient must demonstrate successful completion of each step.
 - Program should be supplemented with a high-repetition, low intensity weight training program focusing on the posterior rotator cuff and scapular musculature.
 - Outcome measures:
 - PSFS: Patient specific functional scale.
 - Quick Dash: Quick disabilities of the arm, shoulder and hand score.
1. Basic menu of program:
- a. Warm-up.
 - b. Stretch.
 - c. 1 set of each exercise prior to ISP.
 - d. ISP.
 - e. 2 sets of each exercise.
 - f. Cryotherapy.